

Research

Sero-epidemiological survey and risk factors associated with brucellosis in dogs in south-western Nigeria

Modupe Comfort Ayoola¹, Akwoba Joseph Ogugua¹, Victor Oluwatoyin Akinseye¹, Tunde Olu Joshua¹, Morenikeji Folusho Banuso², Folashade Julianah Adedoyin¹, Hezekiah Kehinde Adesokan¹, Temidayo Olutayo Omobowale³, John Olusoji Abiola³, Patricia Ihuaku Otuh¹, Helen Oyebukola Nottidge², Emma-Jane Dale⁴, Lorraine Perrett⁴, Andrew Taylor⁴, Judy Stack⁴, Simeon Idowu Babalola Cadmus^{1,&}

¹Department of Veterinary Public Health and Preventive Medicine, Faculty of Veterinary Medicine, University of Ibadan, Ibadan, Nigeria, ²Department of Veterinary Medicine and Surgery, College of Veterinary Medicine, Federal University of Agriculture, Abeokuta, Nigeria, ³Department of Veterinary Medicine, Faculty of Veterinary Medicine, University of Ibadan, Ibadan, Nigeria, ⁴Department of Bacteriology, Animal & Plant Health Agency, United Kingdom

[&]Corresponding author: Cadmus Simeon Idowu Babalola, Department of Veterinary Public Health and Preventive Medicine, Faculty of Veterinary Medicine, University of Ibadan, Ibadan, Nigeria

Key words: Brucellosis, dogs, sero-epidemiology, risk factors, Nigeria

Received: 24/08/2015 - Accepted: 06/12/2015 - Published: 04/02/2016

Abstract

Introduction: In Nigeria, there is limited information on brucellosis particularly in dogs, despite its public health implications. We undertook a sero-epidemiological survey of brucellosis in dogs to determine the prevalence of the disease and associated risk factors for its occurrence in Nigeria. **Methods:** A cross-sectional study was conducted to screen dogs in south-western Nigeria for antibodies to *Brucella sp* using the rapid slide agglutination test (RSA) and Rose Bengal test (RBT), with positive samples confirmed respectively by serum agglutination test (SAT) and competitive enzyme linked immunosorbent assay (cELISA). Data were analyzed with STATA-12. **Results:** From the 739 dog sera tested, 81 (10.96%) were positive by RSA and 94 (12.72%) by RBT; these were corroborated with SAT (4/81; 4.94%) and cELISA (1/94; 1.06%), respectively. Logistic regression identified location (OR=0.04; 95% CI: 0.02-0.09), breed (OR=1.71; 95% CI: 1.34-2.19), age (OR=0.10; 95% CI: 0.04-0.30) and management system (OR=8.51; 95% CI: 1.07-68.05) as risk factors for Brucella infection by RSA. However, location (OR=10.83; 95% CI: 5.48-21.39) and history of infertility (OR=2.62; 95% CI: 1.41-4.84) were identified as risk factors using RBT. **Conclusion:** Given the 10.96% to 12.72% seroprevalence of brucellosis recorded in this study, we advocate control of the disease in dogs, and public health education for those at risk of infection. Again, further studies are required to elucidate the role of dogs in the epidemiology of brucellosis in Nigeria considering the conducive human-animal interface and ecological factors responsible for the transmission of the disease.

Pan African Medical Journal. 2016; 23:29 doi:10.11604/pamj.2016.23.29.7794

This article is available online at: <http://www.panafrican-med-journal.com/content/article/23/29/full/>

© Ayoola Modupe Comfort et al. The Pan African Medical Journal - ISSN 1937-8688. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Introduction

Brucellosis is an infectious disease of global public health importance, with far reaching economic impact since it is associated with reproductive losses in animals [1]. The aetiological agent of the disease is the bacteria of the genus *Brucella*. Generally in canines, *Brucella canis* is the main aetiological agent. It has a ubiquitous distribution and has been reported in the United States, Canada, Central and South America, Tunisia, South Africa, Nigeria, Madagascar, Malaysia, India, Korea, Japan and China among others [2-7]. The organism is not found in New Zealand and Australia [8]. However, brucellosis in dogs can also be caused by *B. abortus*, *B. suis* and *B. melitensis* [9-12] where dogs are in close contact with cattle, sheep, goats and pigs and inadvertently share the same environment. Transmission of *Brucella* infection in dogs occurs via ingestion of contaminated materials or venereal routes [12]. It can also be easily transmitted among dogs reared intensively in breeding kennels or where owners rear two or more dogs. In addition, dogs fed on foetal wastes and raw meats from abattoirs have been reported to be infected with *B. abortus* [13]. The clinical manifestation of the disease in dogs includes abortion, infertility, orchitis, epididymitis and testicular atrophy, among others [14, 15]. Laboratory diagnosis of the disease can be achieved by various serological tests; including the rapid slide agglutination (RSA), indirect fluorescent antibody, serum agglutination test (SAT), agar gel immuno-diffusion assay (AGID) and enzyme linked immunosorbent assay (ELISA). Other tests used are Rose Bengal plate test (RBT), complement fixation tests and fluorescent polarisation assay. However, false positive results may occur due to cross reacting antibodies from other Gram-negative organisms with RBT and RSA [5, 16-18].

While several studies have been conducted on the epidemiology of brucellosis, especially in cattle in Nigeria [19-21], only little is known of the disease among dogs in the country. Whereas, the practice of feeding dogs with foetuses and raw meat from slaughtered cattle coupled with influx of unregistered and suspected brucellosis infected dogs from foreign countries is common. Worse still, there is increasing ownership of dogs by people with poor knowledge of brucellosis, complicated by poor and deplorable health situations in most dog kennels in Nigeria. Based on the aforementioned, it becomes imperative to carry out an epidemiological survey of brucellosis in dogs in Nigeria towards providing empirical data for its control in dogs and humans. To achieve this, we set out to determine the seroprevalence and risk factors associated with brucellosis in dogs in south-western Nigeria.

Methods

Study setting: the study was conducted in Lagos and Ogun States, south-western Nigeria. Lagos State (**Figure 1**) is an administrative division of Nigeria, located in the south-western part of the country and the smallest in land area of Nigeria's 36 states [22]. It is arguably the most economically important state of the country, containing Lagos Division, the nation's largest urban area. Ogun State is another state in south-western Nigeria, located in the north and slightly east to Lagos. Given its contiguous location to Lagos and neighbouring African countries, it also plays vital economic and trans-border activities relating to animal movements and by implication, trans-border diseases. Dogs are reared in both states as pets, and for security as well as for economic purposes.

Study design: We carried out a cross-sectional study. Data from sero-epidemiological survey of dogs presented to major veterinary hospitals/clinics in Lagos and Ogun States were collected between July 2011 and February 2014 for antibodies to *Brucella sp.* In addition, epidemiological data from hunting and stray dogs screened in Ogun State were obtained.

Sample collection and storage: About 5 ml of blood was aseptically collected through the cephalic vein of each sampled dog by a veterinarian. The breed, sex and age of the dogs as well as feed type, and reproduction-related history were obtained and recorded accordingly. The samples were transported to the Tuberculosis and Brucellosis Laboratories of the Department of Veterinary Public Health and Preventive Medicine, University of Ibadan, Nigeria in a cooler. The blood samples were allowed to clot and centrifuged at 3000 x g for five minutes. Serum samples were decanted and stored at -20°C until they were assayed.

Test reagents and procedures: All test reagents used in this study (RSA antigen, RBT antigen, SAT antigen, cELISA kit) were supplied by the Animal and Plant Health Agency (APHA) (Surrey, United Kingdom) and standardized according to the stipulations set by the OIE [23]. Serum samples were examined by RSA, RBT, SAT and cELISA for antibodies to *Brucella sp.* The RSA test and RBT were performed as described by Amin et al. (2012) [24]. Briefly, 30µl of serum sample was mixed with equal volume of antigen on a white enamel plate. The plate was rocked and serum samples that showed agglutination were recorded as sero-positive to *Brucella sp.* Positive samples by RSA and RBT were further respectively analysed using SAT and cELISA as previously described [24, 25]. Data were analyzed using Stata versions 12. Frequencies were generated and Chi square test was used to explore variables potentially associated with *Brucella* infection among dogs. The level of statistical significance was set at $p < 5\%$. Variables significant at 10% on bivariate analysis were entered into the logistic regression model.

Results

Out of 739 dogs screened (Lagos State = 385; Ogun State = 354), an overall seroprevalence of 10.96% and 12.72% were recorded by RSA and RBT out of which, 4.9% (4/81) and 1.1% (1/94) were further supported by SAT and cELISA, respectively. About one third (34.1%) of the dogs screened were Alsatians, more than half (57.6%) were females while two third (65.8%) were adults.

Rapid slide agglutination test

Of 739 serum samples examined by RSA, higher seroprevalence (20.1%) was obtained from Ogun State compared to those from Lagos State (2.6%) (**Table 1**). Breed-specific prevalence was highest among mongrels (38.9%), followed by the Boerboels (8.1%); while the least being the Rottweiler and Alsatian breeds of dogs (1.2%). Also, females had higher seropositivity (11.5%) to antibodies to *Brucella sp.* than the males (10.2%). Age-specific seroprevalence showed that dogs >3 years had higher seroprevalence (13.7%) than those <3 years (5.6%) (**Table 1**). More than one-tenth (12.7%) of dogs with history of infertility were seropositive to antibodies to *Brucella sp.* with only 4.5% from those without infertility. However, a higher seroprevalence of 11.5% was obtained among dogs without history of abortion with only 2.2% from those with previous abortion. Similarly, dogs fed with cow foetus/raw meat had lower seroprevalence (2.8%) than those unexposed to this feed type (13.5%). Seropositivity to antibodies to *Brucella sp.* was also higher among confined (11.2%) than stray

(4.3%) dogs (**Table 1**). The adjusted multivariate logistic regression identified factors like location (OR=0.1; 95% CI: 0.03 - 0.13) and age (OR=6.3; 95% CI: 3.3 - 11.6) to be associated with seropositivity to antibodies to *Brucella sp* (**Table 2**).

Rose Bengal test

From the 739 serum samples examined by RBT, a higher seroprevalence (21.8%) was recorded in Lagos than Ogun (2.8%). The Boerboel breed had the highest breed specific prevalence (21.8%) while the males (13.4%) were more seropositive to antibodies to *Brucella sp* than the females (12.2%) (**Table 3**). Seropositivity was also higher among dogs >3 years (16.2%) than those <3 years (6.0%); while, dogs without history of mating showed higher (14.5%) seroprevalence than those with such history (11.7%). Unexpectedly, lower seroprevalence were obtained among dogs with history of infertility (13.7%) and abortion 8.7%). Similarly, dogs fed with cow fetuses/raw meat (10.2%) and those confined (13.0%) showed higher seropositivity to antibodies to *Brucella sp* (**Table 4**). Overall, location (OR=11.2; 95% CI: 5.7 - 22.1) and infertility (OR=2.6; 95% CI: 1.4 - 4.8) were identified as factors associated with seropositivity to antibodies to *Brucella sp* using the RBT.

Discussion

We report for the first time a large population based brucellosis survey covering diverse dog populations under different settings in Nigeria. The importance of this study is connected with the backdrop of increased dog ownership and very little knowledge about risk factors associated with brucellosis in dogs and its related public health implications in Nigeria. In this study, we employed the use of two *Brucella* antigens namely the standardised RBT antigen and RSA test antigen. The RBT is known to be sensitive to *B. abortus* antibodies, while on the other hand, the RSA is sensitive to *B. canis*. The findings of 10.96% and 12.72% seroprevalence of brucellosis among dogs by RSA and RBT, respectively underscore the importance of this survey in the study area. Importantly, these findings reiterate previous reports [2, 26-28] that brucellosis is endemic in Nigeria. The seroprevalence obtained in this study may be attributed to the fact that brucellosis control programme is non-existent in Nigeria and vaccination of cattle (from locations where dogs could pick up *B. abortus* infection) against brucellosis is not practised [29, 30]. Another source of infection may be infected breeding dogs that shed *Brucella* organism and contaminate dog kennels [31]. Again, the seroprevalence of 10.96% recorded by RSA may not be unconnected with uncontrolled importation of infected dogs from countries with history of *B. canis* in their kennels. These findings portend significant public health implications following the practice of unregulated dog mating without prior screening for brucellosis (a common practice among dog breeders in Nigeria) and associated close contacts between dogs and humans. In addition, poor knowledge of brucellosis [32] coupled with unhygienic practices among dog owners are issues of public health concern that can enhance human infection.

The higher seropositivity recorded by RBT compared with RSA in this study, may be linked with the practice of feeding dogs with cow fetuses/raw meat which is common in the study setting [2]. Again, due to unhygienic practice of disposal of aborted fetuses by herdsman found among West African countries [33], hunting or stray dogs in such environments may consume loads of *Brucella* organisms along with foetal wastes. Thus, this may lead to infection with *B. abortus* which is not a natural pathogen of dogs [10, 26]. Again, we found that seropositivity to *Brucella* infection among dogs

sampled was associated with location of sampling, a finding similar to previous report [2]. As observed, dogs in Lagos were more than nine times more likely to be infected with *Brucella* organism than those in Ogun (using the RBT). This infers that majority of dogs screened in Lagos might be infected with *B. abortus*. This observation may be due to the common practice of importing exotic dogs which are not often screened at the point of entry from neighbouring countries to Lagos. Again since Lagos has a major international airport, the importation corridor therefore makes it easier for more unscreened imported dogs to enter Lagos and thus, the higher population of dog breeders in the state. The implication of this is that Lagos also has more dog breeders most of whom do not keep standard kennel practice; leading to the importation of infected dogs and feeding of dogs with abattoir meat waste (including those originating from cattle with *B. abortus* infection). These assertions are buttressed by Okoh et al. (1978) [34] who isolated *B. canis* from an imported breed of boxer dog in Kano, northern Nigeria; coupled with earlier findings of Cadmus et al. (2011) [2] that associated brucellosis in dogs to feeding of abattoir waste in dogs screened. Therefore, the higher seroprevalence of brucellosis in dogs from Lagos could be due to the more common practice of feeding dogs with foetal waste or raw meat in Lagos than Ogun [2].

Furthermore, our findings identified history of infertility as a factor associated with seropositivity of *Brucella* infection among dogs screened using RBT. Infertility and abortion have been previously reported as major risk factors in the epidemiology of brucellosis among dogs in Ahaz, Iran [6]. Again, infertility is one of the common signs of brucellosis in dogs as well as abortion, failure to conceive, still birth and birth of weak puppies [6, 35]. However, due to the fact that RBT antigen can indiscriminately detect antibodies from other cross reacting organisms, the infertility and abortion identified, to be associated with seropositivity by RBT among dogs screened, may have been due to other organisms. Again, age was identified as a significant factor that plays an important role in the seropositivity of dogs to *Brucella* infection. Our finding showed that adult dogs (>3years) were more than six times more likely to be seropositive to antibodies to *Brucella sp* than the younger ones (OR = 6.3; 95% CI: 3.3 - 11.6). Previous reports similarly indicated that *Brucella* infection in dogs is age-dependent [2, 6, 28]. This could be as a result of longer exposure period with associated higher risks that adult dogs would have been subjected to, an assertion previously corroborated by Kebede et al. (2008) [36].

The breed specific prevalence showed that the mongrel breed is about 30 times more likely to be infected than the Alsatians. This finding may be as a result of inadequate care and attention that dog owners generally give to mongrels (being a local breed and therefore of less commercial value). This attitude may therefore be responsible for the higher exposure to *Brucella* infection. It is noteworthy however, that all canine breeds are equally susceptible to brucellosis [35]. Our finding is similar to that recorded by Cadmus et al. (2011) [2] but in contrast with that recorded in companion dogs in Ahaz, Iran [6], that did not associate the breed of dogs with seropositivity to antibodies to *Brucella sp*. Based on sex of dogs screened, the females were found to be more seropositive than males, though not statistically significant. This observation is similar to the findings by Cadmus et al. (2011) [2] but contrary to the report of Adesiyun et al. (1986) [37]. This may be because a champion stud is more attractive to breeders. Being a source of income to the owner, such stud is usually mated with many females and therefore putting the females at risk of getting infected [2]. Thus, as neither the stud nor bitch is tested, the infected champion stud transmits the disease to many bitches until it becomes apparent that it has become infertile. Similarly, in uncontrolled mating among stray dogs; there is always the alpha male which

mates all the female dogs on heat. Hence, the alpha male could eventually become infected with brucellosis and then transmits same to the females.

Furthermore, results of this study showed that mated animals had lower seroprevalence than the unmated ones, which is in contrast with logical expectation; since brucellosis, can be transmitted during copulation with infected dog. However, since some of the dogs screened may not have been totally confined, it is possible that unwanted mating had taken place without the knowledge of the owners. This observation could therefore have accounted for higher seroprevalence recorded among the "unmated" dogs. Again, confined dogs were found to have higher seroprevalence than the stray dogs, an occurrence which could be attributed to the practice of feeding dogs with fetuses or raw meat. More so, confined dogs may also have higher exposure risk, if at least one of them in the kennel was infected. Despite our findings, this study had its limitations. First, the main screening tests used were the RBT and RSA, while the cELISA and SAT were only used to corroborate results from positive samples. Furthermore, the authors did not carry out bacteriological isolation of *Brucella spp.* This would have provided better insights into the epidemiology of the disease among dogs screened.

Conclusion

This study recorded seroprevalence of 10.96% and 12.72% using RSA and RBT respectively, thus reiterating the fact that brucellosis is prevalent among dogs screened in south-western Nigeria. We also found that location, age and history of infertility are significant factors for infection with *Brucella sp* among dogs. Our findings therefore call for the need to step up routine screening of dogs and public health enlightenment campaigns among dog owners in order to limit the associated hazards on both humans and animals. Finally, control of brucellosis in dogs, will go a long way to prevent zoonotic transmission of the disease, and further avert economic losses associated with adverse reproductive performance in dogs.

Competing interests

The authors declare they have no conflict of interests.

Authors' contributions

Ayoola Modupe Comfort, Ogugua Akwoba Joseph, Akinseye Victor Oluwatoyin, and Adesokan Hezekiah Kehinde participated in sample collection, analysis and wrote the initial draft of the manuscript. Tunde Joshua, Banuso Morenikeji Folusho, Adedoyin Folashade Julianah, Omobowale T.O, Abiola J. O. and Otuh Patricia Ihuaku participated in samples collection and analysis. Nottidge Helen.O, Emma Dale, Lorraine Perrett, Andrew Taylor, Judy Stack, were involved in setting up the study logistics. Cadmus Simeon Idowu Babalola conceived the idea of the study and carried out the final write up of the manuscript. All authors read and approved the final version of the manuscript.

Tables and figures

Table 1: Factors associated with sero-prevalence of brucellosis among dogs in Ogun and Lagos States, south-western Nigeria by RSA

Table 2: Results of logistic regression analysis of factors associated with seroprevalence of brucellosis (with RSA) among dogs

Table 3: Factors associated with sero-prevalence of brucellosis among dogs in Ogun and Lagos States, south-western Nigeria by RBT

Table 4: Results of logistic regression analysis of factors associated with seroprevalence of brucellosis (with RBT) among dogs

Figure 1: Study areas: Lagos and Ogun States (Inset: Nigeria)

References

1. Fernandes ARF, Azevedo SS, Piatti RM, Pinheiro ES, Genovez ME, Azevedo AS, Batista CSA, Alves CJ. *Brucella canis* infection in dogs attended in veterinary clinics from patos, Paraíba state. Brazil Braz J Microbiol. 2011; 142(4):1405-1408. [PubMed](#) | [Google Scholar](#)
2. Cadmus SIB, Adesokan HK, Ajala OO, Odetokun WO, Perrett LL, Stack JA. Seroprevalence of *Brucella abortus* and *B canis* in household dogs in southwestern Nigeria: a preliminary report. Journal of the South African Veterinary Association. 2011; 82(1):56-57. [PubMed](#) | [Google Scholar](#)
3. Ghodasara S, Roy A, Rank DN, Bhandari BB. Identification of *Brucella spp* from animals with reproductive disorders. Buffalo Bulletin. 2010; 29(2):98-108. [PubMed](#) | [Google Scholar](#)
4. Gous TA, Janse van Rensburg, WJ, Gray M, Perrett LL, Brew SD, Young EJ, Whatmore AM, Gers S, Picard J. *Brucella canis* in South Africa. Veterinary Record. 2005; 157(21):668. [PubMed](#) | [Google Scholar](#)
5. Lucero NE, Jacob NO, Ayala SM, Escobar GI, Tuccillo P, Jacques I. Unusual clinical presentation of brucellosis caused by *Brucella canis*. Journal of Medical Microbiology. 2005; 54(5):505-508. [PubMed](#) | [Google Scholar](#)
6. Mosallanejad B, Ghorbanpoor NM, Mohammadian AR. A serological survey on *Brucella canis* in companion dogs in Ahvaz. Iranian J Vet Res. 2009; 10(4): 383 - 386. [PubMed](#) | [Google Scholar](#)
7. Poester FP, Gonçalves VSP, Lage AP. Brucellosis in Brazil. Veterinary Microbiology. 2002; 90(1-4): 55-62. [PubMed](#) | [Google Scholar](#)
8. Larson KL, Spickler AR, Viera S, Dvorak G. Center for Food Security and Public Health. 2012; Iowa State University. USA. [Google Scholar](#)
9. Barr SC, Eilts BE, Roy AF, Miller R. *Brucella suis* biotype 1 infection in a dog. J Am Vet Med Assoc. 1986; 189(6):686-687. [PubMed](#) | [Google Scholar](#)
10. Forbes LB. *Brucella abortus* infection in 14 farm dogs. J Am Vet Med Assoc. 1990; 196:911-916. [PubMed](#) | [Google Scholar](#)
11. Samad MA. Animal Husbandry and Veterinary Science - Vol. 2, 1st Pub, No. 2008. BAU Campus, Mymensingh, Bangladesh. LEP Publication. [Google Scholar](#)

12. Johnson CA, Seguin BE, Davidson AP, Piero FD, Flanders JA, Memon MA, Nicoletti P, Rosenthal RC. Merck Manual for Pet Health, Reproductive disorders of dog. 2011. Kenilworth, N.J., U.S.A. **Google Scholar**
13. Cadmus SIB, Adesokan HK, Adedokun BO, Stack JA. Seroprevalence of bovine brucellosis in trade cattle slaughtered in Ibadan, Nigeria, from 2004-2006. *Journal of South Africa Veterinary Association*. 2010; 81(1):50-53. **PubMed | Google Scholar**
14. Carmichael LE, Greene CE. *Canine brucellosis In: Greene, C.E. (ed) Infectious diseases of the dog and cat - 2nd edition*. 1990. W.B. Saunders, Philadelphia. **Google Scholar**
15. Dunne J, Sehgal K, McMillan A, Perret L. Canine brucellosis in a dog imported into the UK. *Veterinary Record*. 2002; 151(8):247. **PubMed | Google Scholar**
16. Romero C, Gamazo C, Pardo M, López-Goñi I. Specific detection of *Brucella* DNA by PCR. *Journal of Clinical Microbiology*. 1995; 33(3):615-617. **PubMed | Google Scholar**
17. Perrett LL, McGiven JA, Brew SD, Stack JA, John A. Evaluation of Competitive ELISA for Detection of Antibodies to *Brucella* Infection in Domestic Animals. *Croatian Medical Journal*. 2010; 51(4):314-319. **PubMed | Google Scholar**
18. Matero P, Hemmilä H, Tomaso H, Piiparinen H, Rantakokko-Jalava K, Nuotio L, Nikkari S. Rapid field detection assays for *Bacillus anthracis*, *Brucella* spp, *Francisella tularensis* and *Yersinia pestis*, *Clinical Microbiology and Infection*. The Official Publication of the European Society of Clinical Microbiology and Infectious Diseases. 2011; 17(1):34-43. **PubMed | Google Scholar**
19. Adamu NB, Ajogi I. Serological investigation of camels (*Camelus dromedarius*) slaughtered at Kano Municipal Abattoir for evidence of brucellosis. *Tropical Veterinarian*. 1999; 18:45-48. **PubMed | Google Scholar**
20. Cadmus SIB, Ijagbone IF, Oputa HE, Adesokan HK. Serological survey of brucellosis in livestock animals and workers in Ibadan, southwestern, Nigeria. *Afr J Biomed Res*. 2006; 9 (3): 163-168. **PubMed | Google Scholar**
21. Cadmus SIB, Alabi PI, Adesokan HK, Dale EJ, Stack JA. Serological investigation of bovine brucellosis in three cattle production systems in Yewa Division, south-western Nigeria, *Journal of the South African Veterinary Association*. 2013; 84(1):217-222. **Google Scholar**
22. Nigeria Master Web. Nigeria 2006 census figures. Nigeria: Nigeria Master Web; 2006. <http://nigeriamasterweb.com/Nigeria06CensusFigs.htm>. Accessed 16th April 2015. **Google Scholar**
23. OIE. *Manual of diagnostic tests and vaccines for terrestrial animals*, World Organisation for Animal Health, Paris. 2008. **Google Scholar**
24. Amin MM, Ahmed SA, Zaki HM, Ismail RI. 2012, "Serological and molecular studies on the diagnosis of bovine brucellosis". *Nature and Science*. 2012; 10(11):68-76. **PubMed | Google Scholar**
25. Stack J, Perrett L, Brew S, MacMillan A. Competitive ELISA for brucellosis suitable for testing poor quality samples. *Veterinary Record*. 1999; 145(25):735-736. **PubMed | Google Scholar**
26. Baek BK, Lim CW, Rahman MS, Kim CH, Oluoch A, Kakoma I. *Brucella abortus* infection in indigenous Korean dogs. *Can J Vet Res*. 2003; 67(4):312-314. **PubMed | Google Scholar**
27. Sardari K, Kamrani AR, Kazemi KH. The serological survey of *Brucella abortus* and *melitensis* in stray dogs in Mashhad., Iran. *World Small Animal Veterinary Association World Congress Proceedings*. 2003. **Google Scholar**
28. Osinubi MOV, Ajogi I, Ehizibol OD. *Brucella abortus* agglutinin in dogs in Zaria, Nigeria. *Nigeria Veterinary Journal*. 2004; 25(1):35-38. **PubMed | Google Scholar**
29. Rikin UM. Brucellosis of cattle in Nigeria: proposals for a control program under intensive and extensive husbandry systems. 1988. **Google Scholar**
30. Ducrotoy MJ, Bertu WJ, Ocholi RA, Gusi AM, Bryssinckx W, Welburn S, Moriyón I. Brucellosis as an emerging threat in developing economies - Lessons from Nigeria. *PLoS Negl Trop Diseases*. 2014; 8: e3008. **PubMed | Google Scholar**
31. Brower A, Okwumabua O, Massengill C, Muenks Q, Vanderloo P, Duster M, Homb K, Kurth K. Investigation of the spread of *Brucella canis* via the US interstate dog trade. *International Journal of Infectious Diseases*. 2007; 11(5):454-458. **PubMed | Google Scholar**
32. Adesokan HK, Alabi PI, Stack JA, Cadmus SIB. Knowledge and practices related to bovine brucellosis transmission amongst livestock workers in Yewa, south-western Nigeria. *Journal of the South African Veterinary Association*. 2013; 84(1), Art #121: 5 pages. **PubMed | Google Scholar**
33. Unger F, Munstermann S, Goumou A, Apia CN, Konte M, Michaela H. Risk associated with bovine brucellosis in selected study herds and market places in four countries of West Africa (No. Animal Health Working Paper 2). 2003. (p. 37). Banjul, The Gambia. **Google Scholar**
34. Okoh AB, Alexien L, Agbonlahor DE. Brucellosis in dogs in Kano, Nigeria. *Tropical Animal Health and Production*. 1978; 10:210-220. **PubMed | Google Scholar**
35. OIE. *Canine Brucellosis: Brucella canis*. Institute for International Cooperation in Animal Biologics, Iowa State University. 2007. Retrieved from www.cfsph.iastate.edu/IICAB/. **Google Scholar**
36. Kebede T, Ejeta G, Ameni G. Seroprevalence of bovine brucellosis in smallholder farms in central Ethiopia (Wuchale-Jida district). *Revue de Médecine Vétérinaire*. 2008; 159(1):3-9. **PubMed | Google Scholar**
37. Adesiyun AA, Abdullahi SU, Adeyanju JB. Prevalence of *Brucella abortus* and *B canis* antibodies in dogs in Nigeria. *J Small Anim Prod*. 1986; 27: 31-37. **PubMed | Google Scholar**

Table 1: factors associated with sero-prevalence of brucellosis among dogs in Ogun and Lagos States, south-western Nigeria by RSA

Variable	Category	RSA Positive n (%)	Negative n (%)	OR	95% CI	P-value
Location	Ogun State	71(20.1)	283 (79.9)	9.4	4.8 – 18.6	0.00
	Lagos State	10(2.6)	375 (97.4)	1		
Breed	Mongrel	65 (38.9)	102 (61.1)	1		
	Alsatian	3 (1.2)	245 (98.8)	0.01	0.006 – 0.10	0.01
	Boerboel	10 (8.1)	114 (91.9)	0.13	0.06 – 0.30	0.00
	Rottweiler	1 (1.2)	82 (98.8)	0.02	0.002 – 0.14	0.00
	Other	2 (1.7)	115 (98.3)	0.02	0.007 – 0.11	0.00
Sex	Male	32 (10.2)	281 (89.8)	1		
	Female	49 (11.5)	377 (88.5)	1.1	0.7 – 1.8	0.67
Age	<3years	14 (5.6)	237 (94.4)	1		
	>3years	67 (13.7)	421 (96.3)	2.7	1.5 – 4.9	0.00
Mating	Mated	50 (10.8)	413 (89.2)	1		
	Not mated	31 (11.2)	245 (88.8)	1.1	0.6 – 1.7	0.95
Infertility	Fertile	7 (4.5)	148 (95.5)	1		
	Infertile	74 (12.7)	510 (87.3)	3.1	1.3 – 6.8	0.01
Abortion	No	1 (2.2)	45 (97.8)	1		
	Yes	80 (11.5)	613 (88.5)	5.9	0.8 – 43.2	0.08
Fed with fetus/ raw meat	No	5 (2.8)	171 (97.3)			
	Yes	76 (13.5)	487 (86.5)	5.3	2.1 – 13.4	0.00
Management system	Confined	1 (4.3)	22 (95.7)			
	Stray	80 (11.2)	636 (88.8)	2.8	0.4 – 20.8	0.50

Table 2: results of logistic regression analysis of factors associated with seroprevalence of brucellosis (with RSA) among dogs

Variable	Odds ratio	95% CI	p-value
Location			
Ogun State	1		
Lagos State	0.1	0.03 – 0.13	0.00
Age			
<3years	1		
>3years	6.3	3.3 – 11.6	0.00

Table 3: factors associated with sero-prevalence of brucellosis among dogs in Ogun and Lagos States, south-western Nigeria by RBT

Variable	Category	RBT		OR	95% CI	P-value
		Positive n (%)	Negative n (%)			
Location	Ogun State	10 (2.8)	344 (97.2)	1		
	Lagos State	84 (22.1)	301 (77.9)	9.6	4.8 – 18.8	0.00
Breed	Mongrel	8 (5.0)	159 (95.0)	1		
	Alsatian	34 (13.7)	214 (86.3)	3.2	1.4 – 7.0	0.01
	Boerboel	27 (21.8)	97 (78.2)	5.5	2.4 – 12.7	0.00
	Rottweiler	8 (9.6)	75 (90.4)	2.1	0.8 – 5.9	0.23
	Other	17 (14.5)	100 (85.5)	3.4	1.4 – 8.1	0.01
Sex	Male	42 (13.4)	271 (86.6)	1		
	Female	52 (12.2)	374 (87.8)	0.9	0.5 – 1.4	0.71
Age	<3years	15 (6.0)	236 (94.0)	1		
	>3years	79 (16.2)	409 (83.8)	3.1	1.7 – 5.4	0.00
Mating	Mated	54 (11.7)	409 (88.3)	1		
	Not mated	40 (14.5)	236 (85.5)	1.2	0.8 – 2.0	0.32
Infertility	Fertile	14 (9.0)	141 (91.0)	1		
	Infertile	80 (13.7)	504 (86.3)	1.6	0.9 – 2.9	0.16
Abortion	Yes	4 (8.7)	42 (91.3)	1		
	No	90 (13.0)	603 (97.0)	1.6	0.5 – 4.5	0.53
Fed with fetus/ raw meat	Yes	18 (10.2)	158 (89.8)	1		
	No	76 (13.5)	487 (86.5)	1.4	0.8 – 2.4	0.32
Management system	Stray	1 (4.3)	22 (95.7)	1		
	Confined	93 (13.0)	623 (97.0)	3.3	0.4 – 24.7	0.36

Table 4: results of logistic regression analysis of factors associated with seroprevalence of brucellosis (with RBT) among dogs

Variable	Odds ratio	95% CI	p-value
Location			
Ogun State	1		
Lagos State	11.2	5.7 – 22.1	0.00
Infertility			
Fertile	1		
Infertile	2.6	1.4 – 4.8	0.01




Figure 1: Study areas: Lagos and Ogun States (Inset: Nigeria)