

Images in clinical medicine

Primary malignant multiple osteogenic sarcomas: a rare clinical image

 **Sonu Rajwade, Shweta Parwe**

Corresponding author: Shweta Parwe, Department of Panchakarma, Mahatma Gandhi Ayurveda College Hospital and Research Centre, Datta Meghe Institute of Medical Sciences, Wardha, India. drshwetaparve@gmail.com

Received: 31 May 2022 - **Accepted:** 04 Jun 2022 - **Published:** 10 Aug 2022

Keywords: Physical examination, metaphysis, knee joint

Copyright: Sonu Rajwade et al. Pan African Medical Journal (ISSN: 1937-8688). This is an Open Access article distributed under the terms of the Creative Commons Attribution International 4.0 License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Cite this article: Sonu Rajwade et al. Primary malignant multiple osteogenic sarcomas: a rare clinical image. Pan African Medical Journal. 2022;42(263). 10.11604/pamj.2022.42.263.35710

Available online at: <https://www.panafrican-med-journal.com//content/article/42/263/full>

Primary malignant multiple osteogenic sarcomas: a rare clinical image

Sonu Rajwade¹, Shweta Parwe^{1,&}

¹Department of Panchakarma, Mahatma Gandhi Ayurveda College Hospital and Research Centre, Datta Meghe Institute of Medical Sciences, Wardha, India

&Corresponding author

Shweta Parwe, Department of Panchakarma, Mahatma Gandhi Ayurveda College Hospital and Research Centre, Datta Meghe Institute of Medical Sciences, Wardha, India

Image in medicine

We present the case of a 12-year-old male patient, who came to the emergency department with a complaint of pain in bilateral knee joints, pain in bilateral lower legs, multiple swelling over the bilateral knee joint and in the leg region, difficulty in walking, stiffness in the bilateral knee joint since he was one year. There is no medical significant history regarding the disease condition. On physical examination, the swelling was in the region of the metaphysis of the distal end of the bilateral femur and proximal end of the bilateral tibia, skin over the swelling was shiny with tenderness and margins were irregular, on the biopsy and X-ray findings

patient was diagnosed as primary malignant multiple osteogenic sarcomas. The patient was referred to the dermatology department for further management.



Figure 1: primary malignant multiple osteogenic sarcoma of lower limbs