



Research



# Prevalence of modifiable risk factors for cardiovascular disease among school-going children and adolescents in Eldoret, Kenya

#### 🔟 Karani Magutah, 🔟 Grace Wambura Mbuthia, Gilbert Osengo, Dancun Odhiambo, Rebecca Meiring

Corresponding author: Karani Magutah, School of Medicine, Moi University, Eldoret, Kenya. jkarani@cartafrica.org

Received: 05 Dec 2023 - Accepted: 17 Dec 2023 - Published: 04 Mar 2024

Keywords: Cardiovascular disease, risk factors, prevalence, children, adolescents

**Copyright:** Karani Magutah et al. Pan African Medical Journal (ISSN: 1937-8688). This is an Open Access article distributed under the terms of the Creative Commons Attribution International 4.0 License (https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**Cite this article:** Karani Magutah et al. Prevalence of modifiable risk factors for cardiovascular disease among schoolgoing children and adolescents in Eldoret, Kenya. Pan African Medical Journal. 2024;47(100). 10.11604/pamj.2024.47.100.42340

Available online at: https://www.panafrican-med-journal.com/content/article/47/100/full

Prevalence of modifiable risk factors for cardiovascular disease among school-going children and adolescents in Eldoret, Kenya

Karani Magutah<sup>1,&</sup>, Grace Wambura Mbuthia<sup>2</sup>, Gilbert Osengo<sup>1</sup>, Dancun Odhiambo<sup>1</sup>, Rebecca Meiring<sup>3</sup>

<sup>1</sup>School of Medicine, Moi University, Eldoret, Kenya, <sup>2</sup>College of Health Sciences, Jomo Kenyatta University, Juja, Kenya, <sup>3</sup>Department of Exercise Science, University of Auckland, Auckland, New Zealand

#### \*Corresponding author

Karani Magutah, School of Medicine, Moi University, Eldoret, Kenya

# Article 👌



# Abstract

Introduction: Cardiovascular disease (CVD)prevalence in Kenya is rising. Overweight, prehypertension and physical inactivity at younger ages is contributory. These risk factors are inadequately documented among Kenyan children and adolescents, hampering CVD prevention. *Methods:* this cross-sectional study randomly sampled 384 participants from Eldoret, Kenya. After ethical considerations, physical activity was assessed. Body mass index (BMI), Waist-Hip-Ratio (WHR) and Waist-Height-Ratio (WHtR) were determined, and blood pressure (BP) was measured. Results: participants were 14.6±2.7 years, and 62.6% were female. Eight percent had BMI  $\geq$ 25.0 kg/m<sup>2</sup>. Of these, 87% were in secondary schools. Using SBP, 27.9% had CVD risk (42.5% and 20% for males and females ≥13 years and 26.5% and 27% for those <13 years, respectively). For DBP, 12.8% had elevated-to-hypertensive BP (13.2% and 8.3% for males and females ≥13 years and 11.8% and 25.4% for those <13 years, respectively). Combining SBP and DBP, 8.1%, mostly males, had elevated-to-hypertensive BP. Using respective WHR cutoffs of 0.90 and 0.85, 31% (boys) and 15.6% (girls) were at CVD risk. For WHtR, 39.6% of boys were >0.463 cut-off (0.493±0.02) against 32.4% for girls >0.469 cut-off (0.517±0.05). Of these, 52.6% (boys) and 69.7% (girls) were in secondary schools. Overall, 45% of participants were sports-inactive and 77.2% did minimal physical activities. Conclusion: among school-going children and adolescents in Eldoret, Kenya, the prevalence of CVD risk factors was high, especially among boys and in high schools. Large proportions had elevated BP, BMI, WHR and WHtR, and, further, were sedentary, posing a high CVD risk. Lifestyle interventions to mitigate this are urgently needed.

# Introduction

The prevalence of cardiovascular disease (CVD) is increasing in sub-Saharan Africa, a region previously thought to suffer mainly from infectious diseases only. The recently observed trend of increasing sedentary lifestyles in Kenya [1,2] is likely to contribute to the increased prevalence of CVD and associated mortalities in Africa, and will likely result in a significant economic burden to individuals and the country at large [3]. Without intervention, it takes only four years for 20-33% of pre-hypertensive individuals to develop CVD [4]. This statistic is a concern in Kenya as it would be further burden to the economy that already attracts a high per capita cost in the management of CVD [5]. The majority of the Kenyan population is poor, living on <1 USD a day [6].

Much research has identified risk factors for CVD amongst adults [7-9] globally. The prevalence of some risk factors, such as pre-hypertension, is as high as 47% in Kenyan adults (51% for males; 46% for females) [10]. While CVD more commonly occurs in middle-age to older adults, certain modifiable risk factors during childhood and adolescence lead to an increased risk of developing CVD as an adult [11-13]. These risk factors for CVD include overweight and obesity, pre-hypertension, physical inactivity and high sedentary behaviour [11,12,14-16]. Prevalence studies among children and adolescents have however been minimal and the prevalence of modifiable risk factors in these cohorts is not confirmed in Kenya. Recently, one study of primary school children showed 9% had elevated BP while 33% had stage 1 hypertension. In addition, higher BP was associated with higher body mass [17]. While elevated BP and hypertension in children and adolescents continues raising concern not only in both the developed and developing economies [18,19] with a recent systematic review showing prevalence of prehypertension among cohorts in Africa ranging from 2.5-34% [20], we have minimal data from Kenya. In the sub-Saharan region, the prevalence of prehypertension and hypertension in South African primary school children was recently documented ranging 5.2-18.5% [21]. Elevated BP, Prehypertension and hypertension stands at 18.1%, 8.5% 9.6% and respectively in Tanzania [22]. Uncontrolled, these risk factors will



contribute significantly to the development of CVD in adulthood [23].

Interventions such as physical activity (PA) reduce cardiovascular-related morbidity risk and/or modification of the disease for those already affected [24]. When done early in life, such involvement in PA may become a lifestyle carried into adulthood, helping prevent CVD occurrence in those more at risk [15,16,25,26]. School programs screen-time at home have however and contributed to behaviours that are associated with inactivity or sedentariness posing a risk for and the development of enhanced CVD risk factors among school-going children and adolescents [27,28]. We are starting to observe similar trends in Kenya coming at a time when sedentary behavior reduction is now included as an independent factor for consideration in the quest for optimal health [29].

Early screening can help identify at-risk children and adolescents for timely interventions to reduce the occurrence of CVD risk factors. There are realistic and affordable screening approaches available such as the measurements of body composition (BC) and BP as well as methods for the assessment of physical activity and sedentary behaviour through which we are able to find those at a greater risk as well as those already hypertensive. From such measurements we could have an objective documentation of the prevalence of risk factors associated with CVD among children and adolescents in Kenya.

A clear picture of the current prevalence of risk factors associated with the now or future CVD development among Kenyan children and adolescents could help give clarity and, further, evidence needed for designing early interventions necessary to mitigate the problem.

### **Methods**

**Design:** this was a cross-sectional observational study.

**Setting:** participants were drawn from the metropolitan town of Eldoret, western Kenya.

**Study population:** participants were school-going children and adolescents in the upper classes of primary school and high schools in Kenya. This provided a more comparable age bracket since they have similar cut-offs for various variables studied.

Sampling procedure: we studied 384 participants from primary and secondary schools in the metropolitan town of Eldoret, Kenya. The sample size was determined using a CVD risk factors' assumption of 0.5 proportion that would provide largest possible sample, given the the unavailability of Kenyan data for physically inactive/sedentary children and adolescents, or data for other risk factors associated with future development of CVD from Kenya. Participants were drawn from public and private primary and secondary schools. To ease comparisons at analysis, the last four classes of primary school and the four classes of secondary school (as is the system in Kenya) were selected to be invited into the study. The selection for the primary and secondary public and private schools for study was random, picking random papers each with a name of a school from a list of all schools in the metropolis. The numbers studied per school were proportionately allocated based on the student population at each school. To ensure equal chances for participation, selection of the individual participants utilized computer generated random numbers from the class registers.

**Eligibility:** upper primary and secondary schoolgoing children and adolescents within Eldoret were included. Participants excluded were those with physical impairments which could compromise participation in PA as this has been associated with higher CVD risk. These physical challenges are not modifiable risk factors. We also excluded participants with cognitive inabilities that would affect administration of the study tool.





Ethical consideration: this study was approved by the Moi Teaching and Referral Hospital/Moi University Institutional Research Ethics Committee (MTRH/MU IREC) on 22<sup>nd</sup> November 2022 (approval number FAN: 0004291) and obtained research permit from the National Commission for Science, Technology and Innovation (NACOSTI) on 1<sup>st</sup> March 2023 (permit Ref number 839919). After investigators explained the study protocol to participants and answered all related questions, these children and adolescents provided written assent to participate in the study. Since participants were under the legally acceptable age for consenting, their parents/guardians were approached to additionally provide written informed consent for their children to participate.

Assessments: the first participant was recruited on 6<sup>th</sup> March 2023 and engagement with the last participant was on 28<sup>th</sup> April 2023. The global PA questionnaire (GPAQ) was used to assess the level of physical activity and sedentary behaviour. Weight and height were measured using calibrated weighing machine (CAMRY Mechanical scale, BR9012, Shanghai, China) and a tape measure with participants wearing light clothing and no shoes. Waist circumference (WC) was measured with a tape measure, as the diameter horizontally all round just above the iliac crest and hip circumference (HC) as the widest diameter across the buttocks. All measurements were made in centimeters. BP was measured using Omron m<sup>2</sup> Basic (HEM-7120-E) automatic BP monitor (Omron Healthcare Co. Ltd, Kyoto, Japan). Two BP measurements were taken 2 minutes apart following a 5 minutes' rest in a sitting position and the average was recorded

**Data management and analysis:** data were analyzed using STATA v.13 into proportions, means and standard deviation. Missing data was skipped for individual participants, causing a change in specific variables' sample size "n" accordingly. Analyses for PA engagement was based on proportions that were either active or sedentary. All variables were thereafter re-analyzed based on applicable cut-offs to identify participants' risk levels. Participants with values above their set cut-offs for all or either of Waist-Height-Ratio (WHtR) (0.463 for boys and 0.469 for girls) [30-32], Waist-Hip-Ratio (WHR) ( $\geq$ 0.90 (male) and  $\geq$ 0.85 (female) [33], body mass index (BMI) (a 95 percentile of adult values applied amongst <13 year old and a similar percentile used for BMI categorization for all participants' age range) [34,35] and BP ( $\geq$ 120/80 for ages  $\geq$ 13 years and a 95 percentile for < 13 year old) [35] were considered at risk for CVD.

### **Results**

**Bio-demographic characteristics:** there were 145 (37.4%) male and 243 (62.6%) female participants. The average age of participants was 14.6 $\pm$ 2.7 years. Those  $\geq$ 13 years (n=285) had mean age 15.6 $\pm$ 2.2 years while those <13 years (n=99) had 11.4 $\pm$ 0.8 mean age. Thirty-four percent (33.8%), 13.1%, 30.5% and 22.6% were drawn from public secondary, private secondary, private primary and public primary schools respectively. The class distributions for participants ranged from 8.5% for the lowest representation (grade 5) to 20.5% for the highest (class 8) using the Kenyan education system.

Blood pressure-related risk factors: eight percent (7.6%, n=29) of the participants had hypertensive pressure ≥130 mmHg (≥124 mmHg for <13 years old) using SBP while 18.5% had prehypertensive ranges 120-129 mmHg (114-123mmHg for <13 years old). For males and females respectively, 25% (n=35) and 14.8% (n=36) were prehypertensive while 9.3% (n=13) and 6.6% (n=16) had hypertensive SBPs. Using DBP, 7.3% (n=28) of participants were in elevated BP range of 80-89 mmHg (≥76-85 mmHg) for <13 years old) and 1.3% (n=5) were already hypertensive. For males and females respectively, 7.9% (n=11) and 7% (n=17) had prehypertensive pressures, with 0.7% (n=1) and 1% (n=4) having hypertensive DBPs ≥90 mmHg ( $\geq$ 86 mmHg for < 13 years old). Combining both SBP and DBP, 8.1% of participants (n=31) accounted for by a 9.3% (n=13) among males and 7.4% among females (n=18) had elevated to



hypertensive BPs. A chi-squared test showed that overall, males had higher SBP compared to females ( $\chi^2$  = 9.81; p = 0.007) and that those from private schools had higher SBP when compared to those from the public ( $\chi^2$  = 15.54; p = <0.001). The various means for the cardiovascular and body composition variables for the 144 males and 243 females are shown in Table 1.

Body composition-related risk factors: six percent (6.2%) of participants (N=387) had a BMI of 25-<30 and 1.6% were at ≥30. Disaggregating to males (N=144) and females (N=243), 5.6% versus 6.6% were in the 25-<30 BMI range respectively. While 2.5% females had BMI ≥30, no males had such. Being female was associated with higher BMI ( $\chi^2$  = 11.26; p = 0.01). Eighty-six point six percent (86.6%) of those with BMI ≥25.0 were in secondary schools and which was associated with higher BMI  $(\chi^2 = 102.54; p < 0.001)$ , and 53% came from public schools. The mean (SD) WC for males was 72.9±9.2 centimeters and 71.3±10.5 centimeters for females. Using proposed ≥84.6cm for boys and ≥72cm for girls (given as 90<sup>th</sup> percentile of African adult WC value ≥94 and 80cm in males and females respectively) as cut-offs for WC corresponding with high CVD risk [36], 32.4% of participants (11% of boys, WC 89.8± 4.9; 45.1% of girls, WC 79.6±9.1) had higher-than-recommended values. The WHR was 0.86±0.06 and 0.79±0.07 for boys and girls respectively. Using respective WHR cut-offs of 0.90 and 0.85 for males and females, 30.8% of boys were at CVD risk (WHR 0.93±0.02) while the girls at risk were 15.6% (mean 0.89±0.03). For the boys (n=45), 42.2% were in secondary schools but for the girls (n=38), half each were from primary and secondary schools. For WHtR, it was 0.45±0.04 and 0.45±0.06 for boys and girls. For the boys, 39.6% had their WHtR above 0.463 cut-off (mean 0.493±0.02). Distribution of those with WHtR above cut-off (n=57) was 29.8% from private secondary schools, 26.3% from private primary schools, 22.8% from public secondary schools, and, 21.1% from public primary schools. Overall, 52.6% came from secondary schools while the rest were in primary. For the girls, those above respective 0.469 cut-off

were 32.4% (mean 0.517±0.05). Distribution of the girls at-risk using WHtR (n=79) had 16.5% from private secondary schools, 21.5% from private primary schools, 53.2% from public secondary schools and 8.9% from public primary schools. Overall, 69.6% of girls with at-risk WHtR were in high schools compared to 30.4% in primary schools. About 45% of the participants did not participate in any form of planned exercise activities (46.8% among males, 44.2% among females). Outside planned exercises, 77.2% (78.9% male; 76.3% female) did not participate in appreciable daily moderate or vigorous intensity PA at home or school. Distribution across schools for this and related cardiovascular and body composition variables is shown in Table 2.

## **Discussion**

Participants in the current study were in midteens. This age has been observed to have significant modifiable CVD risk factors that include overweight and obesity, pre-hypertension, physical inactivity and high sedentary behaviours. These risk factors are more common later during the middle-ages and increase through older adulthood. When they start early in teen ages, the likelihood of enhanced CVD risk in adulthood is increased [11,12,14].

Almost 1 in 4 participants had elevated-tohypertensive SBPs for their ages. In every 3 males, one had elevated-to-hypertensive SBPs and there were almost twice as many affected as the females. Using DBP, about 8% were in elevated-tohypertensive ranges and the males were still more affected. Combining both SBP and DBP also showed that males are much more affected compared to the females in our current study. A similar trend where males are more affected than females has been shown before among adults aged  $\geq$ 18 years in Kenya, and, more comparatively age-wise, in neighbouring Uganda and Tanzania [10,37,38]. While previously adult data showed up to 47% of Kenyan adults are prehypertensive, the current study shows slightly lower proportions although it involved younger





participants. The at-risk proportions in the current study are within ranges found regionally in the sub-Saharan Africa and the larger continent [20,37-39]. With previous observation that up to 30% of children and adolescents aged ≥12 years who have prehypertension develop CVD later in life [15,16], the high proportions with prehypertension in the current study may need to be addressed. There were mixed results for the mean SBP and DBP values for the males and females but higher SBP mainly affected the males. While recent works locally and regionally, just like the current study show fairly high proportions with prehypertension and hypertensive BPs among primary school children [17,20,22,37-39], what the current study adds is that those in private schools had higher proportions with elevated-to-hypertensive SBPs. Further, the current study adds that those in secondary schools are more affected with about 1 in 3 having prehypertensive-to-hypertensive BPs, and, more than twice as affected as their counterparts in respective primary school category. The differences in proportions were not marked for DBP, although those from high schools were higher. The findings compares to those of a study done in Congo that showed increased odds of prehypertensive among school children in secondary and private schools compared to those in primary and public schools respectively [39].

Although most participants had BMI within acceptable ranges, four in every ten, and mostly boys, were underweight. Importantly given the focus of the study, appreciable proportions had overweight-to-obese BMIs, and the females were most affected. Participants from secondary schools were especially affected, probably because they were older, and public schools had a slight edge over the private. Similar findings were documented in a meta-analysis of prevalence of overweight and obesity among African primary school learners where overweight and obesity estimates were higher in private compared to public schools. This could point to the difference in socio-economic status and thus lifestyle behaviors of the families where these children come from.

For WC, while the mean WC appeared within acceptable range for both sexes based on age [36], a third of all participants had higher-thanrecommended values. Girls had four times more proportions affected, with just under a half of all females having higher than acceptable WC. It is safe to say that overall, by use of WC alone, 1 in 3 participants from the current study had central obesity and therefore enhanced CVD risk. Using respective WHR cut-offs however showed males having twice as much proportions affected compared to females and about half of such boys and girls were in both school categories. The WHtR, also showed males as having larger proportions at risk, with 2 in every 5 having WHtR >0.463. The majority of these were drawn from both private and secondary schools. For females, 1 in 3 had values >0.469 cut-off but here, although those from public secondary schools contributed the bulk overall, their proportions per school category remained, like for all other BC variables, lower compared to private schools. For all BC variables, proportions for those at higher risk were much more from high than primary schools, and, mostly, among the private.

While exercise and PA are known to reduce CVD risks [24], almost half of the current study participants did not participate in any form of planned moderate and/or vigorous intensity exercises. The participation was poorer among boys compared to girls. Similarly for PA at home or in school, the males fared slightly worse for those with minimal-to-no PA involvement. Overall, almost 4 in 5 participants were sedentary using this criterion, an exact observation first made a decade ago but among older adults [1], and, also, more recently in a study following COVID-19 associated lifestyle changes that showed little progress has occurred in exercise and PA participation in Eldoret, Kenya [2]. If something is not done and these children and adolescents take a sedentary lifestyle of minimal PA involvement into their adult lives, their CVD risk could end up higher. Distribution across schools showed that higher proportions of children and adolescents from private schools did not participate in





exercises and PA, and that those in high schools for both categories had lesser engagement in such planned exercises. It has previously been observed that school programs and screen-time contribute the inactivity being observed among school children and adolescents [27,28], and although the current study did not seek this association, it is likely this contributes to the minimal exercises and PA participation observed here.

Limitations: we used self-reports for PA and exercise data and this subjective data may not only have introduced recall bias but also compromised quality. Given that this was a crosssectional study and, further, data was collected during school sessions made this inevitable. Data for the BC were collected by 2 research assistants and although they had been trained together, there may have been individual differences in the accuracy of the measurements taken. Further, BP measurements were done once-off in a single session. These may have affected the results.

### Conclusion

Based on the BC, BP and PA screening used in the current study, school-going children and adolescents in Eldoret, Kenya had high prevalence of CVD risk-factors. This was especially higher among boys and in high schools. Large proportions had elevated BPs, BMI, WHR and WHtR, and, further, were sedentary, posing high CVD risk. Lifestyle interventions to mitigate this public health concern and that target on children and adolescents are urgently needed.

#### What is known about this topic

- The prevalence of cardiovascular disease is increasing in Kenya and the larger sub-Saharan Africa region;
- Prevalence of risk factors for CVD is high among Kenyan adults;
- Without intervention, it takes only four years for 1 in 4 individuals at risk for CVD to develop it.

#### What this study adds

- This study found that a quarter of children and adolescents in Eldoret, Kenya have elevated-to-hypertensive SBPs for their ages; like has been found for sex distribution among older individuals, the current study found that more Kenyan boys than girls have elevated-to-hypertensive blood pressure, and, using systolic BP alone, twice as many boys as girls have this CVD risk factor;
- The study also found higher proportions of children and adolescents from private schools with elevated-to-hypertensive SBPs, and those in secondary schools as most affected;
- The study adds that using the various body composition measurements, 1 in 3 children and adolescents in western Kenya have central obesity.

## **Competing interests**

The authors declare no competing interests.

## Funding

Karani Magutah was supported by the Consortium for Advanced Research Training in Africa (CARTA). CARTA is jointly led by the African Population and Health Research Center and the University of the Witwatersrand and funded by the Carnegie Corporation of New York (Grant No. G-19-57145), Sida (Grant No: 54100113), Uppsala Monitoring Center, Norwegian Agency for Development Cooperation (Norad), and by the Wellcome Trust [reference no. 107768/Z/15/Z] and the UK Foreign, Commonwealth and Development Office, with support from the Developing Excellence in Leadership, Training and Science in Africa (DELTAS Africa) programme. The statements made and views expressed are solely the responsibility of the Fellow.



# Authors' contributions

Karani Magutah conceptualized the idea, designed the study, sourced for funding and wrote the first draft. Additionally, he together with Grace Mbuthia, analyzed the data. Gilbert Osengo and Duncan Odhiambo collected all data. All stated authors, together with Rebeca Meiring, participated in refining final manuscript. All authors read and approved the final version of the manuscript.

## Acknowledgments

The authors acknowledge the various school heads and teachers who allowed and facilitated data collection from their respective schools.

# **Tables**

**Table 1**: cardiovascular and body compositionmeans and proportions

**Table 2**: percentage of participants not meetingexpected levels in various variables per schoolcategory

### References

- Nambakai JE, Kamau J, Amusa LO, Goon DT, Andaje M. Factors influencing participation in physical exercise by the elderly in Eldoret West District, Kenya. African Journal for Physical, Health Education, Recreation and Dance (AJPHERD), 2011;17(3): 462-472. Google Scholar
- Magutah K, Mbuthia G. Physical activity engagement in Eldoret, Kenya, during COVID-19 pandemic. PLOS Glob Public Health. 2(4): e0000339. PubMed | Google Scholar
- Oyando R, Njoroge M, Nguhiu P, Kirui F, Mbui J, Sigilai A *et al*.Patient costs of hypertension care in public health care facilities in Kenya. Int J Health Plann Manage. 2019;34(2): e1166e1178. PubMed| Google Scholar

- Vasan RS, Larson MG, Leip EP, Kannel WB, Levy D. Assessment of frequency of progression to hypertension in nonhypertensive participants in the Framingham Heart Study: a cohort study. Lancet. 2001;358(9294): 1682-6. PubMed| Google Scholar
- Oti SO, van de Vijver S, Gomez GB, Agyemang C, Egondo T, Kyobutungi C *et al.* Outcomes and costs of implementing a communitybased intervention for hypertension in an urban slum in Kenya. Bull World Health Organ. 2016;94(7): 501-9. PubMed | Google Scholar
- World Bank Report. Kenya. Accessed 12<sup>th</sup> February 2020.
- Roth GA, Mensah GA, Johnson CO, Addolorato G, Ammirati E, Baddour LM *et al.* Global Burden of Cardiovascular Diseases Writing Group (2020). Global Burden of Cardiovascular Diseases and Risk Factors, 1990-2019: Update From the GBD 2019 Study. J Am Coll Cardiol. 2020 Dec 22;76(25): 2982-3021. PubMed | Google Scholar
- Ruan Y, Guo Y, Zheng Y, Huang Z, Sun S, Kowal P et al. Cardiovascular disease (CVD) and associated risk factors among older adults in six low-and middle-income countries: results from SAGE Wave 1. BMC Public Health. 2018;18(1): 77. PubMed| Google Scholar
- Choukem SP, Tochie JN, Sibetcheu AT, Nansseu JR, Hamilton-Shield JP. Overweight/obesity and associated cardiovascular risk factors in sub-Saharan African children and adolescents: a scoping review. Int J Pediatr Endocrinol. 2020;2020: 6. PubMed | Google Scholar
- Ogola E, Yonga G, Njau K. A14369 High Burden of Prehypertension in Kenya: Results from the Healthy Heart Africa (HHA) program. Journal of Hypertension. 2018;36(): e330. Google Scholar
- Weihrauch-Blühe S, Schwarz P, Klusmann JH. Childhood obesity: increased risk for cardiometabolic disease and cancer in adulthood. Metabolism. 2019 Mar;92: 147-152. PubMed | Google Scholar



- Jacobs DR Jr, Woo JG, Sinaiko AR, Daniels SR, Ikonen J, Juonala M *et al.* Childhood Cardiovascular Risk Factors and Adult Cardiovascular Events. N Engl J Med. 2022;386(20): 1877-1888. PubMed| Google Scholar
- Sommer A, Twig G. The Impact of Childhood and Adolescent Obesity on Cardiovascular Risk in Adulthood: a Systematic Review. Curr Diab Rep. 2018 Aug 30;18(10): 91. PubMed| Google Scholar
- 14. Han M, Li Q, Liu L, Zhang D, Ren Y, Zhao Y et al. Prehypertension and risk of cardiovascular diseases: a meta-analysis of 47 cohort studies. J Hypertens. 2019;37(12): 2325-2332.
  PubMed | Google Scholar
- Redwine KM, Daniels SR. Pre-Hypertension in Adolescents: Risk and Progression. J Clin Hypertens (Greenwich). 2012;14(6): 360-364.
   PubMed | Google Scholar
- Redwine KM, Falkner B. Progression of prehypertension to hypertension in adolescents. Curr Hypertens Rep. 2012;14(6): 619-25. PubMed | Google Scholar
- Gewa CA, Onyango AC, Opiyo RO, Gittelsohn J, Cheskin LJ. Patterns and predictors of elevated blood pressure and hypertension among primary school children in urban Kenya. J Hypertens. 2022;40(12): 2513-2520.
   PubMed | Google Scholar
- Riley M, Hernandez AK, Kuznia AL. High Blood Pressure in Children and Adolescents. Am Fam Physician. 2018;98(8): 486-494. PubMed| Google Scholar
- Redjala O, Ahmed MS, Cherifi M, Smati L, Benhassine F, Baghriche M *et al.* Children hypertension in Northern Africa. Am J Cardiovasc Dis. 2021;11(2): 222-230.
   PubMed | Google Scholar
- Malik KS, Adoubi KA, Kouame J, Coulibaly M, Tiade ML, Oga S *et al.* Prevalence and Risks Factors of Prehypertension in Africa: A Systematic Review. Annals of global health. 2022;88(1): 13. PubMed | Google Scholar

- 21. Gomwe H, Seekoe E, Lyoka P, Marange CS. Blood pressure profile of primary school children in Eastern Cape province, South Africa: prevalence and risk factors. Ann Glob Health. 2022 Mar 1;88(1): 13. PubMed| Google Scholar
- 22. Sungwa EE, Kibona SE, Dika HI, Laisser RM, Gemuhay HM, Kabalimu TK *et al.* Prevalence and factors that are associated with elevated blood pressure among primary school children in Mwanza Region, Tanzania. Pan Afr Med J. 2020 Nov 30: 37: 283. PubMed| Google Scholar
- 23. Seeman T, Hamdani G, Mitsnefes M. Hypertensive crisis in children and adolescents. Pediatr Nephrol. 2019;34(12): 2523-2537. PubMed | Google Scholar
- 24. Zhuang Z, Gao M, Yang R, Li N, Liu Z, Cao W *et al.* Association of physical activity, sedentary behaviours and sleep duration with cardiovascular diseases and lipid profiles: a Mendelian randomization analysis. Lipids Health Dis. 2020;19(1): 86. PubMed| Google Scholar
- Niiranen TJ, Larson MG, McCabe EL, Xanthakis V, Vasan RS, Cheng S. Prognosis of Prehypertension Without Progression to Hypertension. Circulation. 2017;136(13): 1262-1264. PubMed | Google Scholar
- Egan BM and Stevens FS. Prehypertension---prevalence, health risks, and management strategies. Nat Rev Cardiol. 2015;12(5): 289-300. PubMed | Google Scholar
- 27. Grao-Cruces A, Sánchez-Oliva D, Padilla-Moledo C, Izquierdo-Gómez R, Cabanas-Sánchez V, Castro-Piñero J. Changes in the school and non-school sedentary time in youth: The UP&DOWN longitudinal study. J Sports Sci. 2020;38(7): 780-786. PubMed| Google Scholar
- 28. Pierce JB, Kershaw KN, Kiefe CI, Jacobs DR Jr, Sidney S, Merkin SS *et al.* Association of Childhood Psychosocial Environment With 30-Year Cardiovascular Disease Incidence and Mortality in Middle Age. J Am Heart Assoc. 2020 May 5;9(9): e015326. PubMed| Google Scholar



- 29. Bull FC, Al-Ansari SS, Biddle S, Borodulin K, Buman MP, Cardon G *et al.* World Health Organization 2020 guidelines on physical activity and sedentary behaviour. Br J Sports Med. 2020 Dec;54(24): 1451-1462. **PubMed**| **Google Scholar**
- Kruger HS, Faber M, Schutte AE, Ellis SM. A proposed cutoff point of waist-to-height ratio for metabolic risk in African township adolescents. Nutrition. 2013;29(3): 502-507.
   PubMed | Google Scholar
- 31. Matsha TE, Kengne AP, Yako YY, Hon GM, Hassan MS, Erasmus RT. Optimal waist-toheight ratio values for cardiometabolic risk screening in an ethnically diverse sample of South African urban and rural school boys and girls. PloS One. 2013;8(8): e71133. PubMed| Google Scholar
- 32. Jiang Y, Dou YL, Xiong F, Zhang L, Zhu GH, Wu T *et al.* Waist-to-height ratio remains an accurate and practical way of identifying cardiometabolic risks in children and adolescents. Acta Paediatr. 2018 Mar 22. Epub ahead of print. **PubMed** | **Google Scholar**
- 33. WHO Waist circumference and waist-hip ratio: Report of a WHO expert consultation Geneva 8-11. 2008. Google Scholar
- 34. Centers for Disease Control and Prevention A SAS Program for the 2000 CDC Growth Charts (ages 0 to <20 years) Division of Nutrition Physical Activity and Obesity, N.C.f. and Disease Control and Prevention Atlanta GA USA 2014.

- 35. Gidding SS, Whelton PK, Carey RM, Flynn J, Kaelber DC, Baker-Smith C. Aligning Adult and Pediatric Blood Pressure Guidelines. Hypertension. 2019;73(5): 938-943. PubMed| Google Scholar
- 36. Xi B, Zong X, Kelishadi R, Litwin M, Hong YM, Poh BK et al. International Waist Circumference Percentile Cutoffs for Central Obesity in Children and Adolescents Aged 6 to 18 Years. J Clin Endocrinol Metab. 2020 Apr 1;105(4): e1569-83. PubMed| Google Scholar
- 37. Katamba G, Agaba DC, Migisha R, Namaganda A, Namayanja R, Turyakira E. Prevalence of hypertension in relation to anthropometric indices among secondary adolescents in Mbarara, Southwestern Uganda. Ital J Pediatr. 2020 Jun 1;46(1): 76. PubMed| Google Scholar
- 38. Nsanya MK, Kavishe BK, Katende D, Mosha N, Hansen C, Nsubuga RN *et al.* Prevalence of high blood pressure and associated factors among adolescents and young people in Tanzania and Uganda. J Clin Hypertens (Greenwich). 2019;21(4): 470-478. PubMed| Google Scholar
- 39. Ellenga Mbolla BF, Okoko AR, Mabiala Babela JR, Ekouya Bowassa G, Gombet TR, Kimbally-Kaky SG *et al.* Prehypertension and hypertension among schoolchildren in Brazzaville, Congo. Int J Hypertens. 2014: 2014: 803690. **PubMed** | Google Scholar

<b>Fable 1:</b> cardiovascular and body composition means and proportions							
Variable				% > cut-off			
	Males	Females	Overall	Males	Females	overall	
BMI (kg/m²)	18.9±3.3	20.4±4.1	19.8±3.9	5.6	9.1	7.8	
SBP (mmHg)	112.3±13.6	108.9±12.0	110.1±12.7	38.5	21.8	27.9	
≥13 years	115.7±12.0	109.3±11.9	111.7±12.3	42.5	20.0	28.3	
< 13 years	101.9±13.3	107.7±12.1	105.6±12.8	26.5	27.0	26.8	
DBP (mmHg)	67.2±8.4	68.8±7.7	68.2±8.0	12.9	12.8	12.8	
≥13 years	67.8±8.6	68.6±7.3	68.3±7.8	13.2	8.3	10.1	
< 13 years	65.3±7.6	69.7±8.8	68.1±8.6	11.8	25.4	20.8	
WC (cm)	72.9±9.2	71.3±10.5	*	11	45.1	32.4	
WHR	0.86±0.06	0.79±0.07	*	30.8	15.6	21.3	
WHtR	0.45±0.04	0.45±0.06	*	39.6	32.4	35.1	

BMI, body mass index; kg/m<sup>2</sup>, kilogram per metre squared; SBP, systolic blood pressure; DBP, diastolic blood pressure; mmHg, millimeters of mercury; WC, waist circumference; cm, centimetres; WHR, waist hip ratio; WHtR, waist height ratio. Data are mean ± SD unless otherwise indicated. For SBP and DBP, males ≥13 and <13 years n=106 and 34 respectively, and, for the females, n = 180 (age ≥13 years) and 63 (age <13 years) respectively. \*Recommended cut-offs differ for boys and girls

Table 2: percentage of participants not meeting expected levels in various variables per school category								
Variable	% of total in PuHS	% of total in	% of total in PuPS	% of total in PrPS				
	(n=131)	PrHS (n=51)	(n=119)	(n=88)				
BMI (>25)	12.2	19.6	0	3.4				
<b>SBP</b> (≥ 13 years, ≥120mmHg; <13-	29.7	37.2	22.7	23.8				
year-old, ≥114 mmHg)								
<b>DBP</b> (≥13 years, ≥80mmHg; <13	10.9	7.8	11.7	13.6				
years, ≥76 mmHg)								
wc								
Males (≥84.6 cm)	17.1	25.9	0	5.4				
Females (≥72 cm)	63.9	75	11.7	38.1				
WHR								
Males (≥0.9)	20	40.7	46.4	23.2				
Females (≥0.85)	12.4	29.2	13.3	14.3				
WHtR								
Males (≥0.463)	40	63	46.4	26.8				
Females (≥0.469)	43.3	54.2	11.7	27.0				
Exercises (non-participating proportion)	44.4	69.4	20.6	54.2				
<b>PA</b> (non-participating proportion)	49.2	83.7	93.2	92.4				
PuHS, Public high school: PrHS, priv	vate high school: Pu	PS, public prima	ary school: PrPS, priv	ate primary school				

PuHS, Public high school; PrHS, private high school; PuPS, public primary school; PrPS, private primary school; BMI, body mass index; SBP, systolic blood pressure; DBP, diastolic blood pressure; WC, waist circumference; WHR, waist hip ratio; WHtR, waist height ratio; PA, physical activity.