

Images in clinical medicine



Delayed sequelae of proximal tibial fracture presenting with advanced arthritis managed by total knee arthroplasty

Anjali Nawkhare, Mitushi Deshmukh

Corresponding author: Anjali Nawkhare, Department of Musculoskeletal Physiotherapy, Ravi Nair Physiotherapy College, Datta Meghe Institute of Higher Education and Research, Sawangi (Meghe), Wardha, Maharashtra, India. anjalinawkhare@gmail.com

Received: 13 Sep 2025 - Accepted: 25 Sep 2025 - Published: 03 Nov 2025

Keywords: Total knee arthroplasty, sequela of proximal tibial fracture, tibial plateau, orthopaedic surgery

Funding: This work received no specific grant from any funding agency in the public, commercial, or non-profit sectors.

Copyright: Anjali Nawkhare et al. Pan African Medical Journal (ISSN: 1937-8688). This is an Open Access article distributed under the terms of the Creative Commons Attribution International 4.0 License (https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Cite this article: Anjali Nawkhare et al. Delayed sequelae of proximal tibial fracture presenting with advanced arthritis managed by total knee arthroplasty. Pan African Medical Journal. 2025;52(92). 10.11604/pamj.2025.52.92.49358

Available online at: https://www.panafrican-med-journal.com//content/article/52/92/full

Delayed sequelae of proximal tibial fracture presenting with advanced arthritis managed by total knee arthroplasty

Anjali Nawkhare^{1,&}, Mitushi Deshmukh¹

¹Department of Musculoskeletal Physiotherapy, Ravi Nair Physiotherapy College, Datta Meghe Institute of Higher Education and Research, Sawangi (Meghe), Wardha, Maharashtra, India

[&]Corresponding author

Anjali Nawkhare, Department of Musculoskeletal Physiotherapy, Ravi Nair Physiotherapy College, Datta Meghe Institute of Higher Education and Research, Sawangi (Meghe), Wardha, Maharashtra, India

Article 3



Image in medicine

Six (6) years after receiving surgery for a compound grade 3B fracture of the proximal right tibia, a 45-year-old man with a history of bronchial asthma came with chronic right knee pain and limited mobility. Magnetic resonance imaging (MRI) showed that the tibiofemoral and patellofemoral joints had diminishing articular cartilage and signs of a chronic cleavage fracture affecting both tibial plateaus. Along with a degenerative tear of the posterior horn,

degenerative alterations were observed in both horns and the lateral meniscus body. Radiographs showed subchondral sclerosis and narrowing of the joint space, which are post-traumatic osteoarthritic alterations. The patient underwent successful total knee arthroplasty (TKA), leading to marked improvement in joint function and pain relief. The patient had a successful functional recovery following a total right knee replacement. This case emphasizes the long-term degenerative effects of high-grade open tibial plateau fractures and how total knee replacement can help restore joint function in these complex situations.



Figure 1: A) magnetic resonance imaging of the right knee demonstrating chronic post-traumatic changes, including thinning of the articular cartilage, degenerative tear of the lateral meniscus, and evidence of an old cleavage fracture involving both tibial plateaus; B) radiographs showing postoperative anteroposterior and lateral views of the right knee showing a well-aligned total knee replacement (TKR) implant with femoral component and tibial stem extension; the presence of cerclage wires and screws indicates prior fixation, consistent with previous management of a complex tibial plateau fracture