

Images in clinical medicine

Intraoperative stapled anastomosis following small bowel perforation: a clinical image

Sachin Bharshankar, Ruchira Ankar

Corresponding author: Ruchira Ankar, Department of Medical Surgical Nursing, Smt. Radhikabai Meghe Memorial College of Nursing, Datta Meghe Institute of Higher Education and Research, Wardha, Maharashtra, India. ruchira3110@gmail.com

Received: 14 Oct 2025 - **Accepted:** 04 Nov 2025 - **Published:** 06 Feb 2026

Keywords: Exploratory laparotomy, abdominal closure, Linear gastrointestinal stapler

Funding: This work received no specific grant from any funding agency in the public, commercial, or non-profit sectors.

Copyright: Sachin Bharshankar et al. Pan African Medical Journal (ISSN: 1937-8688). This is an Open Access article distributed under the terms of the Creative Commons Attribution International 4.0 License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Cite this article: Sachin Bharshankar et al. Intraoperative stapled anastomosis following small bowel perforation: a clinical image. Pan African Medical Journal. 2026;53(62). 10.11604/pamj.2026.53.62.49784

Available online at: <https://www.panafrican-med-journal.com//content/article/53/62/full>

Intraoperative stapled anastomosis following small bowel perforation: a clinical image

Sachin Bharshankar¹, Ruchira Ankar^{1,&}

¹Department of Medical Surgical Nursing, Smt. Radhikabai Meghe Memorial College of Nursing, Datta Meghe Institute of Higher Education and Research, Wardha, Maharashtra, India

&Corresponding author

Ruchira Ankar, Department of Medical Surgical Nursing, Smt. Radhikabai Meghe Memorial College of Nursing, Datta Meghe Institute of Higher Education and Research, Wardha, Maharashtra, India

Image in medicine

A 50-year-old male patient presented with acute onset abdominal pain associated with vomiting, progressive abdominal distension, and failure to pass flatus and feces; and peritonitis due to distal ileal perforation. Emergency exploratory laparotomy with bowel resection and stapled side-to-side anastomosis was performed, followed by thorough peritoneal lavage. There was a history of a long-standing abdominal wall swelling that became painful and irreducible. On examination, the patient was toxic with signs of intestinal obstruction and localized peritonism. Routine demographic identifiers are withheld to maintain

anonymity. An emergency exploratory laparotomy was undertaken following clinical and radiological suspicion of acute intra-abdominal pathology. Intraoperatively, a diseased segment of small bowel. Hemodynamic parameters stabilized within the first 24 hours, and a gradual resolution of abdominal pain and distension was the short-term outcome. Among the medium-term outcomes were: successful tolerance of oral feeding by day 4-5; normal bowel movements restoration;

complete wound healing without dehiscence; and no anastomotic leak oriented tra-abdominal abscess. The diagnostic approach included: acute abdomen with features of bowel obstruction and strangulation; laboratory investigations: leukocytosis and metabolic derangement suggestive of sepsis; and radiological assessment: abdominal imaging suggestive of obstructed hernia.




Figure 1: intraoperative view showing stapled side-to-side small bowel anastomosis after resection of the perforated ileal segment